

EUROPEAN PATENT APPLICATION

(21) Application number: 89306453.5

(22) Date of filing: 26.06.89

(51) Int. Cl. 4: C07D 487/04 , C07D 495/04 ,  
 C07D 498/04 , C07D 513/04 ,  
 A61K 31/505 , C07D 231/14 ,  
 C07D 333/38 , C07D 285/10 ,  
 C07D 271/10 , //(C07D487/04,  
 239:00,231:00),(C07D495/04,  
 333:00,239:00),(C07D498/04,  
 271:00,239:00),(C07D513/04,  
 285:00,239:00)

Claims for the following Contracting States: ES  
 + GR

The title of the invention has been amended  
 (Guidelines for Examination in the EPO, A-III,  
 7.3).

(30) Priority: 01.07.88 GB 8815716  
 01.07.88 GB 8815717  
 01.07.88 GB 8815718

(43) Date of publication of application:  
 03.01.90 Bulletin 90/01

(54) Designated Contracting States:  
 AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: SMITH KLINE & FRENCH


LABORATORIES LIMITED  
 Mundells  
 Welwyn Garden City Hertfordshire, AL7  
 1EY(GB)

(72) Inventor: Coates, William John  
 3 Rocks Hill Welwyn Garden City  
 Hertfordshire AL8 1EU(GB)  
 Inventor: Rawlings, Derek Anthony,  
 75 Stanmore Road Stevenage  
 Hertfordshire SG1 3QE(GB)

(73) Representative: Waters, David Martin, Dr. et al  
 Smith Kline & French Laboratories Ltd.  
 Patent Department Mundells  
 Welwyn Garden City Hertfordshire AL7  
 1EY(GB)

(57) Condensed pyrimidine derivatives, process and intermediates for their preparation and  
 pharmaceutical compositions containing them.

(57) Compounds of the formula (1):




(1)


and pharmaceutically acceptable salts are described wherein




is a ring of sub-formula (a), (b) or (c):



(a)



(b)



(c),

X is oxygen or sulphur, and


R<sup>1</sup> is C<sub>1</sub>-<sub>6</sub>alkyl, C<sub>2</sub>-<sub>6</sub>alkenyl, C<sub>3</sub>-<sub>6</sub>cycloalkyl, C<sub>1</sub>-<sub>4</sub>alkyl, or C<sub>1</sub>-<sub>4</sub>alkyl substituted by 1 to 6 fluoro groups.

Their use as medicaments, pharmaceutical compositions containing them and processes for their preparation are also described.

# CHEMICAL COMPOUNDS

The present invention relates to fused pyrimidine derivatives, processes for their preparation, intermediates in their preparation, their use as therapeutic agents and to pharmaceutical compositions containing them. The compounds of this invention are inhibitors of a calmodulin insensitive cyclic GMP phosphodiesterase and are of use in combatting such conditions where such inhibition is thought to be beneficial. They are bronchodilators and are therefore of use in combatting chronic reversible obstructive lung diseases such as asthma and bronchitis. Furthermore the compounds of this invention are vasodilators and are therefore of value in combatting angina, hypertension and congestive heart failure.


Accordingly the present invention provides compounds of the formula (1) :




and pharmaceutically acceptable salts thereof, wherein




is a ring of sub-formula (a), (b) or (c):



(a)



(b)



(c) ,

X is oxygen or sulphur, and

R¹ is C₁-alkyl, C₂-alkenyl, C₃-cycloalkyl, C₁-alkyl substituted by 1 to 6 fluoro groups.

Suitably R¹ is C₂-alkyl for example ethyl, n-propyl, isopropyl, butyl, isobutyl or pentyl.

Suitably R¹ is C₃-alkenyl for example allyl, butenyl or pentenyl.

Suitably R¹ is cyclopropylmethyl.

Examples of C₁-alkyl substituted by 1 to 6 fluoro groups include -CF₃, -CH₂CF₃ or -CF₂CHFCF₃.

Preferably R¹ is n-propyl.

Suitably



is group of sub-formula (a) thus forming a pyrazolo[3,4-d]pyrimidine ring system.

Suitably



5 is a group of sub-formula (b) thus forming a thieno[2,3-d]pyrimidine ring system.  
Suitably



10

is a group of sub-formula (c) and X is oxygen thus forming a [1,2,5]oxadiazolo[3,4-d]pyrimidine ring system.  
Suitably

15



20 is a group of sub-formula (c) and X is sulphur thus forming a [1,2,5]thiadiazolo[3,4-d]pyrimidine ring system.

Particular compounds of this invention are:

- 6-(2-propoxyphenyl)pyrazolo[3,4-d]pyrimidin-4(5H)-one,
- 2-(2-propoxyphenyl)thieno[2,3-d]pyrimidin-4(3H)-one,
- 2-(2-propoxyphenyl)[1,2,5]oxadiazolo[3,4-d]pyrimidin-4(3H)-one, or
- 25 2-(2-propoxyphenyl)[1,2,5]thiadiazolo[3,4-d]pyrimidin-4(3H)-one

or pharmaceutically acceptable salts thereof.

This invention covers all tautomeric and optical isomeric forms of compounds of formula (1)

Compounds of the formula (1) may form pharmaceutically acceptable salts with metal ions, alkali metals for example sodium and potassium, or with an ammonium ion.

30 In order to use a compound of the formula (1) or a pharmaceutically acceptable salt thereof in the treatment of humans and other mammals it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

Compounds of formula (1) and their pharmaceutically acceptable salts may be administered in standard manner for the treatment of the indicated diseases, for example orally, sublingually, parenterally, transdermally, rectally, via inhalation or via buccal administration.

35 Compounds of formula (1) and their pharmaceutically acceptable salts which are active when given orally or via buccal administration can be formulated appropriately in dosage forms such as liquids, syrups, tablets, capsules and lozenges. An oral liquid formulation will generally consist of a suspension or solution of the compound or salt in a liquid carrier for example, ethanol, glycerine or water with a flavouring or colouring agent. Where the composition is in the form of a tablet, any pharmaceutical carrier routinely used for preparing solid formulations may be used. Examples of such carriers include starch, celluloses, lactose, sucrose and magnesium stearate. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example using the aforementioned carriers in a hard gelatin capsule shell.

45 Where the composition is in the form of a soft gelatin shell capsule any pharmaceutical carrier routinely used for preparing dispersions or suspensions may be considered, for example aqueous gums, celluloses, silicates or oils and are incorporated in a soft gelatin capsule shell.

Typical parenteral compositions consist of a solution or suspension of the compound or salt in a sterile aqueous or non-aqueous carrier optionally containing a parenterally acceptable oil, or solubilising agent, for example polyethylene glycol, polyvinylpyrrolidone, 2-pyrrolidone, cyclodextrin, lecithin, arachis oil, or sesame oil.

50 A typical suppository formulation comprises a compound of formula (1) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent, for example polymeric glycols, gelatins, cocoa-butter or low melting vegetable waxes or fats or their synthetic analogues.

55 Typical transdermal formulations comprise a conventional aqueous or non-aqueous vehicle, for example a cream, ointment, lotion or paste or are in the form of a medicated plaster, patch or membrane.

Typical compositions for inhalation are in the form of a solution, suspension or emulsion that may be

administered in the form of an aerosol using a conventional propellant such as dichlorodifluoromethane or trichlorofluoromethane, or are in the form of a powder for insufflation.

Preferably the composition is in unit dosage form, for example a tablet, capsule or metered aerosol dose, so that the patient may administer to himself a single dose.

5 Each dosage unit for oral administration contains suitably from 0.001 mg/Kg to 30 mg/Kg, and preferably from 0.005 mg/Kg to 15 mg/Kg, and each dosage unit for parenteral administration contains suitably from 0.001 mg/Kg to 10 mg/Kg, of a compound of formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base.

10 The daily dosage regimen for oral administration is suitably about 0.001 mg/Kg to 120 mg/Kg, of a compound of formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base. The daily dosage regimen for parenteral administration is suitably about 0.001 mg/Kg to 40 mg/Kg, for example about 0.005 mg/Kg to 10 mg/Kg, of a compound of the formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base. The active ingredient may be administered as required for example from 1 to 8 times a day or by infusion. The compositions of the invention are bronchodilators and are useful  
15 in chronic reversible obstructive lung disease for example asthma and bronchitis. The compositions of the present invention also have vasodilator activity and are of use in the treatment of angina, hypertension and congestive heart failure. Such conditions can be treated by administration orally, sublingually, topically, rectally, parenterally or by inhalation. For administration by inhalation dosages are controlled by a valve, are administered as required and for an adult are conveniently in the range 0.1-5.0 mg of a compound of the  
20 formula (1) or a pharmaceutically acceptable salt thereof.


The compounds of this invention may be co-administered with other pharmaceutically active compounds, for example in combination, concurrently or sequentially. Conveniently the compounds of this invention and the other active compound or compounds are formulated in a pharmaceutical composition. Examples of compounds which may be included in pharmaceutical compositions with the compounds of the  
25 formula (1) are bronchodilators such as sympathomimetic amines for example isoprenaline, isoetharine, salbutamol, phenylephrine and ephedrine or xanthine derivatives for example theophylline and aminophylline, anti-allergic agents for example disodium cromoglycate, histamine H<sub>1</sub>-antagonists, vasodilators for example hydralazine, angiotensin converting enzyme inhibitors for example captopril, or agents for example isosorbide nitrate, glyceryl trinitrate and pentaerythritol tetranitrate, anti-  
30 example quinidine, procainamide and lignocaine, calcium antagonists for example verapamil, diuretics such as thiazides and related compounds for example bendroflumazide, chlorothalidone, hydrochlorothiazide, and other diuretics for example frusemide and triamterene, and sedatives for example nitrazepam, flurazepam and diazepam.

In another aspect the present invention provides a process for the preparation of compounds of the  
35 formula (1) or pharmaceutically acceptable salts thereof, which process comprises:

a) for compounds wherein




is a group of sub-formula (a), reacting a compound of the formula (2):



50 wherein R<sup>1</sup> is as hereinbefore defined with a compound of the formula (3):


55



10 wherein L<sup>1</sup> is C<sub>1</sub>-alkoxy or amino;  
b) for compounds wherein



is a group of sub-formula (a) or (b), cyclising a compound of the formula (4):




wherein R<sup>1</sup> is as hereinbefore defined and



35 is a group of sub-formula (a) or (b) as hereinbefore defined;  
c) for compounds wherein




is a group of sub-formula (c) and X is oxygen, cyclising a compound of the formula (5):



55 wherein R<sup>1</sup> is as hereinbefore defined with an oxidising agent;  
d) for compounds wherein

A

5 is a group of sub-formula (c) and X is sulphur, reacting a compound of the formula (6) :



wherein R¹ is as hereinbefore defined with thionyl chloride or a chemical equivalent thereof; and thereafter optionally forming a pharmaceutically acceptable salt.

20 Suitably an acid addition salt of a compound of the formula (2), for example the methanesulphonate or hydrochloride, is reacted with an acid addition salt of a compound of the formula (3), for example the sulphate or hydrochloride, in the presence of a suitable base such as sodium acetate in the absence of a solvent or in a suitable solvent such as a C₁-12 alcohol, pyridine or N-methylpyrrolidin-2-one at an elevated temperature, for example 50-250 °C.


25 Suitably a compound of the formula (4) is cyclised by heating at an elevated temperature, for example 50-150 °C, in the presence of an acid or a base in a suitable solvent such as aqueous C₁-12 alcohols, water, toluene, a halohydrocarbon or acetonitrile. Conveniently a compound of the formula (4) is cyclised by heating in aqueous base such as sodium hydroxide at the reflux temperature of the reaction mixture.

Suitably a compound of the formula (5) is reacted with an oxidising agent in a solvent such as acid at a moderate temperature e.g. from 0-80 °C, conveniently at ambient temperature. A suitable oxidising agent is lead tetraacetate.

30 Suitably an acid addition salt of compound of the formula (6), for example the sulphate or hydrochloride, is reacted with excess thionyl chloride or a chemical equivalent thereof in the absence of a solvent or in the presence of a solvent such as a halohydrocarbon at an elevated temperature for example from 50-200 °C, conveniently at the reflux temperature of the reaction mixture. By a chemical equivalent thereof is meant a reagent that will react with a compound of the formula (6) in a similar manner to thionyl chloride to afford a compound of the formula (1) wherein X is sulphur, for example N-thionylaniline (J. Org. Chem., 29, 2135 (1964)).

The compounds of the formulae (2), (5) and (6) and acid addition salts thereof are known or preparable in conventional manner from US Patent 3819631.

40 A compound of the formula (4) can be prepared by reaction of 3-amino-4-pyrazolecarboxamide or 2-aminothiophene-3-carboxamide with a compound of the formula (7):



wherein R¹ is as hereinbefore defined and L² is halo.

Suitably L² is chloro or bromo. Suitably a compound of the formula (7) is reacted with 3-amino-4-pyrazole-carboxamide at ambient or elevated temperature e.g. 50-100 °C in a suitable solvent such as toluene, acetonitrile or a halohydrocarbon e.g. chloroform or dichloromethane, optionally in the presence of a base such as pyridine or triethylamine, to form a compound of the formula (4) which may be cyclised in situ or may be isolated and thereafter cyclised as hereinbefore described.

Pharmaceutically acceptable base addition salts of the compounds of the formula (1) may be prepared by standard methods, for example by reacting a solution of the compound of the formula (1) with a solution

of the base.

The following biological test method, data and Examples serve to illustrate this invention.

## 5 Bronchodilatation - In vivo

Male guinea-pigs of the Dunkin Hartley strain (500 - 600g) were anaesthetised with Sagatal (pentobarbital sodium) (60 mg/kg). Airway resistance was measured using a modification of the classical Konzett-Rossler technique (J. Pharm. Methods, 13, 309-315, 1985). U46619 (9,11-methanoepoxy-PGH<sub>2</sub>) was infused i.v. at a rate of 2.5 nmol/min, this produced a steady state of bronchoconstriction (approximately 120% increase from basal airway resistance). The compound under test was administered by i.v. bolus injection, and the subsequent peak inhibition of bronchoconstriction recorded.

The dose of compound required to reduce the U46619-induced bronchoconstriction by 50% is given as the BD<sub>50</sub>. This result demonstrates in vivo anti-bronchoconstrictor activity.

Compound of Example	BD <sub>50</sub> (μmol/kg)
1	5.13
2	2.62
4	4.51

### 25 Example 1

## 30 6-(2-Propoxyphenyl)pyrazolo[3,4-d]pyrimidin-4(5H)-one

A stirred mixture of 2-propoxybenzamidinium methane-sulphonate (1.0 g), 3-amino-4-pyrazolocarboxamide sulphate (0.5 g) and sodium acetate (0.82 g) was heated in an oil bath (180 °C) for one hour. The reaction mixture was dissolved in hot 2 Normal sodium hydroxide (10 ml) and the solution was treated with charcoal and neutralised with glacial acetic acid. A precipitate was collected, washed with water and eluted from a silica column with chloroform. The combined fractions containing product were evaporated to dryness to afford the title compound, 148 mg, m.p. 182-183.5 °C.

### 40 Example 2

## 45 2-(2-Propoxyphenyl)thieno[2,3-d]pyrimidin-4(3H)-one

a) A solution of 2-propoxybenzoyl chloride (0.99 g) in acetonitrile (7.5 ml) was added dropwise over 5 minutes to a cooled (0 °C), stirred mixture of 2-aminothiophene-3-carboxamide (0.71 g) and triethylamine (0.51 g) in acetonitrile (7.5 ml). The reaction mixture was stirred for one hour while being allowed to warm to ambient temperature and was then allowed to stand overnight. Acetonitrile was removed under reduced pressure and the residue was washed with water and recrystallised from ethanol-water to afford 2-(2-propoxybenzamido)thiophene-3-carboxamide, 0.91 g, m.p. 176.5-178.5 °C.

b) A stirred mixture of 2-(2-propoxybenzamido)thiophene-3-carboxamide (0.90 g) and pyridine (1 ml) in 2 Normal sodium hydroxide (25 ml) was heated under reflux for 2 hours. The cooled reaction mixture was neutralised with concentrated hydrochloric acid to afford a precipitate and the resulting mixture was extracted with chloroform (3 x 25 ml). The combined extracts were washed with water (20 ml) and brine (20 ml), dried (magnesium sulphate) and evaporated under reduced pressure to afford a crude product which was recrystallised from ethanol-water to afford the title compound, 0.41 g, m.p. 115-117 °C.

### Example 3

#### 5 2-(2-Propoxyphenyl)[1,2,5]oxadiazolo[3,4-d]pyrimidin-4(3H)-one

Lead tetraacetate (1 g) was added portionwise over 5 minutes to a stirred solution of 4-amino-5-nitroso-2-(2-propoxyphenyl)pyrimidin-6-one (0.52 g) in acetic acid (10 ml) under nitrogen. The reaction mixture was stirred for 1.5 hours and then allowed to stand overnight. A precipitate was collected, washed with water, dissolved in hot methanol and treated with charcoal. The methanolic solution was filtered through diatomaceous earth and the filtrate was evaporated under reduced pressure to afford a residue which was recrystallised from methanol to afford the title compound, 0.19 g, m.p. 163-164° C.

### 15 Example 4

#### 20 2-(2-Propoxyphenyl)[1,2,5]thiadiazolo[3,4-d]pyrimidin-4(3H)-one

A stirred solution of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one sulphate (0.72 g) in thionyl chloride (25 ml) was heated under reflux for 2 hours. The cooled reaction mixture was evaporated under reduced pressure and the residue was washed with water and recrystallised from methanol-water and then from methanol to afford the title compound, 0.38 g, m.p. 157.5-159.5° C.

### 25 Example 5

30 Pharmaceutical compositions for oral administration are prepared by combining the following :

	% w/w		
	0.5	3.0	7.14
2-(2-Propoxyphenyl)thieno[2,3-d]pyrimidin-4(3H)-one	30.45	88.2	64.41
2% w/w Soya lecithin in soya bean oil	9.05	8.8	8.45
Hydrogenated vegetable shortening and beeswax			


35 The formulations are then filled into individual soft gelatin capsules.

### 40 Example 6

45 A pharmaceutical composition for parenteral administration is prepared by dissolving the title compound of Example 4 (0.02 g) in polyethylene glycol 300 (25 ml) with heating. This solution is then diluted with water for injections Ph. Eur. (to 100 ml). The solution is then sterilised by filtration through a 0.22 micron membrana filter and sealed in sterile containers.

### 50 Claims

1. A compound of the formula (1) :




(1)


or a pharmaceutically acceptable salt thereof, wherein




is a ring of sub-formula (a), (b) or (c):



(a)



(b)



(c),

X is oxygen or sulphur, and

- R¹ is C₁-₆ alkyl, C₂-₆ alkenyl, C₃-₆ cycloalkyl, C₁-₆ alkyl, or C₁-₆ alkyl substituted by 1 to 6 fluorine.
2. A compound according to claim 1 wherein R¹ is C₂-₆ alkyl.
  3. A compound according to claim 1 wherein R¹ is n-propyl.
  4. A compound according to any one of claims 1 to 3 where



is a group of sub-formula (a).

5. A compound according to any one of claims 1 to 3 where



is a group of sub-formula (b).

6. A compound according to any one of claims 1 to 3 where



is a group of sub-formula (c) and X is oxygen.

7. A compound according to any one of claims 1 to 3 where

A

is a group of sub-formula (c) and X is sulphur.

8. A compound according to claim 1 which is :

6-(2-propoxyphenyl)pyrazolo[3,4-d]pyrimidin-4(5H)-one,

2-(2-propoxyphenyl)thieno[2,3-d]pyrimidin-4(3H)-one,

10 2-(2-propoxyphenyl)[1,2,5]oxadiazolo[3,4-d]pyrimidin-4(3H)-one, or  
2-(2-propoxyphenyl)[1,2,5]thiadiazolo[3,4-d]pyrimidin-4(3H)-one,

or a pharmaceutically acceptable salt thereof.

9. A compound according to any one of claims 1 to 8 for use as a medicament.

10. A pharmaceutical composition which comprises a compound according to any one of claims 1 to 8  
and a pharmaceutically acceptable carrier.


15 11. A process for preparing a compound of the formula (1) or a pharmaceutically acceptable salt thereof  
as defined in claim 1 which comprises :

a) for compounds wherein

20

A


25 is a group of sub-formula (a), reacting a compound of the formula (2):



30

wherein  $R^1$  is as defined in claim 1 with a compound of the formula (3):

35



40

wherein  $L^1$  is  $C_1-4$ alkoxy or amino;

45


b) for compounds wherein

A

50

is a group of sub-formula (a) or (b), cyclising a compound of the formula (4):

55



10 wherein R¹ is as hereinbefore defined and




is a group of sub-formula (a) or (b) as defined in claim 1;

c) for compounds wherein



25 is a group of sub-formula (c) and X is oxygen, cyclising a compound of the formula (5) :




35 wherein R¹ is as hereinbefore defined with an oxidising agent;

40 d) for compounds wherein



is a group of sub-formula (c) and X is sulphur, reacting a compound of the formula (6) :




wherein R<sup>1</sup> is as hereinbefore defined with thionyl chloride or a chemical equivalent thereof; and thereafter optionally forming a pharmaceutically acceptable salt.

12. A compound of the formula (4) as defined in claim 11.

5 Claims for the following Contracting State: ES


1. A process for preparing a compound of the formula (1):



20 or a pharmaceutically acceptable salt thereof, wherein




is a ring of sub-formula (a), (b) or (c):




X is oxygen or sulphur, and  
R<sup>1</sup> is C<sub>1</sub>-6 alkyl, C<sub>2</sub>-6 alkenyl, C<sub>3</sub>-6 cycloalkyl, C<sub>1</sub>-4 alkyl, or C<sub>1</sub>-4 alkyl substituted by 1 to 6 fluoro groups,  
which process comprises:  
40 a) or compounds wherein



is a group of sub-formula (a), reacting a compound of the formula (2):




wherein R<sup>1</sup> is as hereinbefore defined with a compound of the formula (3):



wherein L<sup>1</sup> is C<sub>1</sub>-alkoxy or amino;  
 10 b) for compounds wherein



15 is a group of sub-formula (a) or (b), cyclising a compound of the formula (4) :




wherein R<sup>1</sup> is as hereinbefore defined and



35 is a group of sub-formula (a) or (b) as defined in claim 1;  
 c) for compounds wherein




is a group of sub-formula (c) and X is oxygen, cyclising a compound of the formula (5) :



55 wherein R<sup>1</sup> is as hereinbefore defined with an oxidising agent;  
 d) for compounds wherein

5 is a group of sub-formula (c) and X is sulphur, reacting a compound of the formula (6) :



wherein R<sup>1</sup> is as hereinbefore defined with thionyl chloride or a chemical equivalent thereof;  
and thereafter optionally forming a pharmaceutically acceptable salt.

- 20
2. A process according to claim 1 for preparing a compound wherein R<sup>1</sup> is C<sub>2-5</sub> alkyl.
  3. A process according to claim 1 for preparing a compound wherein R<sup>1</sup> is n-propyl.
  4. A process according to any one of claims 1 to 3 for preparing a compound wherein

25

is a group of sub-formula (a).

- 30
5. A process according to any one of claims 1 to 3 for preparing a compound wherein

35

is a group of sub-formula (b).

6. A process according to any one of claims 1 to 3 for preparing a compound wherein

40

is a group of sub-formula (c) and X is oxygen.

- 45
7. A process according to any one of claims 1 to 3 for preparing a compound wherein

50

is a group of sub-formula (c) and X is sulphur.

8. A process according to claim 1 for preparing a compound which is :

6-(2-propoxyphenyl)pyrazolo[3,4-d]pyrimidin-4(5H)-one,

2-(2-propoxyphenyl)thieno[2,3-d]pyrimidin-4(3H)-one,

55 2-(2-propoxyphenyl)[1,2,5]oxadiazolo[3,4-d]pyrimidin-4(3H)-one, or


2-(2-propoxyphenyl)[1,2,5]thiadiazolo[3,4-d]pyrimidin-4(3H)-one,

or a pharmaceutically acceptable salt thereof.

9. A process for preparing a pharmaceutical composition which comprises bringing into association a

compound of the formula (1) as defined in any one of claims 1 to 8 and a pharmaceutically acceptable carrier.


10. A process for preparing a compound of the formula (4) as defined in claim 1 which comprises reacting 3-amino-4-pyrazolocarboxamide or 2-aminothiophene-3-carboxamide with a compound of the formula (7):



wherein R<sup>1</sup> is as defined in claim 1 and L<sup>2</sup> is halo.

Claims for the following Contracting State : GR




1. A process for preparing a compound of the formula (1):



or a pharmaceutically acceptable salt thereof, wherein




is a ring of sub-formula (a), (b) or (c):




X is oxygen or sulphur, and  
R<sup>1</sup> is C<sub>1</sub>-<sub>6</sub>alkyl, C<sub>2</sub>-<sub>6</sub>alkenyl, C<sub>3</sub>-<sub>5</sub>cycloalkyl, C<sub>1</sub>-<sub>4</sub>alkyl, or C<sub>1</sub>-<sub>4</sub>alkyl substituted by 1 to 6 fluoro groups,  
which process comprises:  
a) for compounds wherein



is a group of sub-formula (a), reacting a compound of the formula (2):




wherein R<sup>1</sup> is as hereinbefore defined with a compound of the formula (3) :



wherein L<sup>1</sup> is C<sub>1</sub>-alkoxy or amino;  
b) for compounds wherein



is a group of sub-formula (a) or (b), cyclising a compound of the formula (4) :




wherein R<sup>1</sup> is as hereinbefore defined and



is a group of sub-formula (a) or (b) as defined in claim 1;  
c) for compounds wherein



is a group of sub-formula (c) and X is oxygen, cyclising a compound of the formula (5) :




(5)

wherein R¹ is as hereinbefore defined with an oxidising agent;  
d) for compounds wherein



is a group of sub-formula (c) and X is sulphur, reacting a compound of the formula (6) :



(6)

wherein R¹ is as hereinbefore defined with thionyl chloride or a chemical equivalent thereof;  
and thereafter optionally forming a pharmaceutically acceptable salt.

2. A process according to claim 1 for preparing a compound wherein R¹ is C₂-5 alkyl.
3. A process according to claim 1 for preparing a compound wherein R¹ is a propyl.
4. A process according to any one of claims 1 to 3 for preparing a compound wherein



is a group of sub-formula (a).

5. A process according to any one of claims 1 to 3 for preparing a compound wherein



is a group of sub-formula (b).

6. A process according to any one of claims 1 to 3 for preparing a compound wherein



is a group of sub-formula (c) and X is oxygen.

7. A process according to any one of claims 1 to 3 for preparing a compound wherein

5 is a group of sub-formula (c) and X is sulphur.

8. A process according to claim 1 for preparing a compound which is :

6-(2-propoxyphenyl)pyrazolo[3,4-d]pyrimidin-4(5H)-one,

2-(2-propoxyphenyl)thieno[2,3-d]pyrimidin-4(3H)-one,

10 2-(2-propoxyphenyl)[1,2,5]oxadiazolo[3,4-d]pyrimidin-4(3H)-one, or

2-(2-propoxyphenyl)[1,2,5]thiadiazolo[3,4-d]pyrimidin-4(3H)-one,

or a pharmaceutically acceptable salt thereof.

9. A process for preparing a pharmaceutical composition which comprises bringing into association a compound of the formula (1) as defined in any one of claims 1 to 8 and a pharmaceutically acceptable carrier.

15 10. A compound of the formula (4) as defined in claim 1.

20

25

30

35

40

45

50

55